

Impact of gravity waves on the middle atmosphere of Mars: a non-orographic gravity wave parameterization based on Global Climate modeling and MCS observations

G. Gilli^{1,2}, F. Forget², A. Spiga², T. Navarro³, E. Millour², L. Montabone^{2,4},

A. Kleinböhl⁵, D. M. Kass⁵, D. J. McCleese⁶, J. T. Schofield⁵

¹Instituto de Astrofísica e Ciências do Espaço (IA), Lisbon, Portugal

²Laboratoire de Météorologie Dynamique (LMD/IPSL), Centre National de la Recherche Scientifique, Sorbonne Université, École Normale Supérieure, École Polytechnique, Paris, France

³Department of Earth, Planetary, and Space Sciences, University of California, Los Angeles, CA 90095-1567, USA

⁴Space Science Institute, Boulder, Colorado, USA

⁵Jet Propulsion Laboratory, California Institute of Technology, Pasadena, California, USA

⁶California Institute of Technology, Pasadena, California, USA

Contents of this file

Figure S1.

Introduction This supporting information provides a Figure showing an example of zonal mean wind (m/s) simulated by the LMD-MGCM at standard resolution 64x48x32 (as in the paper), compared with higher vertical resolution 64x48x54 runs with the LMD-MGCM. The figure shows that increasing the vertical resolution from 6-7 km (i.e. 32 vertical levels) to 2-3 km (i.e. 54 vertical levels) does not have a significant impact

on the GW drag, demonstrating that using a refined resolution is not critical in our study. Similarly to other parameterizations, the non-orographic GW parameterization implemented in our model does not strongly depend on the vertical resolution.




Figure S1. Left panels: Simulated LMD-MGCM zonal wind performed using the standard resolution 64x48x32 (upper panel) for early northern hemisphere winter ($L_s=270^\circ$) with the non-orographic GW scheme include, as in Figure 4 (panel c), together with higher vertical resolution 64x48x54 (lower panel) for the same L_s . Right panels: corresponding (daily averaged) GW drag on the zonal wind.